SERICULTURE EXTENSION OFFICER (RPC) IN THE DEPARTMENT OF SERICULTURE

Examination held on 20-08-2023 (AN)

Specific Paper (Paper - 2) (Subject Code: 545)

KEY CHART

Question Booklet Series - 'S'

Total Marks: 300

3 Marks for each question

Question Number	Key Answer	Question Number	Key Answer	Question Number	Key Answer	Question Number	Key Answer
1	1	26	1	51	2	76	3
2	1	27	4	52	3	77	2
3	3	28	1	53	1	78	3
4	2	29	4	54	2	79	1
5	4	30	2	55	2	80	4
6	3	31	2	56	2	81	3
7	3	32	3	57	2	82	1
8	1	33	1	58	Th	83	1
9	2	34	1	59	~(1)	84	3
10	3	35	3	60	Ci	85	4
11	3	36	3	61	* 3	86	4
12	1	37	4	62	2	87	3
13	4	38	3	63	3	88	4
14	3	39	2	64	4	89	1
15	2	40	3	65	3	90	4
16	2	41	3	66	3	91	2
17	4	42	C4	67	2	92	2
18	4	43	3	68	2	93	1
19	3	44	2	69	4	94	2
20	4	45	3	70	2	95	2
21	1	46	2	71	1	96	4
22	2	47	2	72	4	97	1
23	2	48	3	73	4	98	3
24	2	49	2	74	4	99	2
25	4	50	1	75	3	100	2

Note: Every question for which wrong answer has been given by the candidate, ¼ th (0.25) of the marks assigned for that question will be deducted.

Sd/-Controller of Examination Karnataka Public Service Commission Bengaluru.

1.	Th	e mussel silk is produced by :	4.	The	gene identified for expression of
	(1)	Pinna squamosa		"Gol	den yellow coloured cocoon" is :
ŧ	(2)	Samia cynthia ricini		(1)	Y
	(3)	Pachypasa otus		(2)	C
	(4)	Nephila madagascariensis		(3) (4)	+ ^y I
3.	in	e following insect pest plays a key role cross infectivity of pebrine during tworm rearing: Mulberry leaf roller Thrips Mealy bug Mites Multiplication Farms are also	5.	Give State is sh cross State grow In light	n below are two statements: ment A: The larval span of females orter than that of males in reciprocal ies. ment B: The females of hybrids at a faster rate. ght of above statements, choose the appropriate answer. Statement A is correct but Statement B is incorrect >
	calle	ed:		(2).	Statement B is correct but
	(1)	P4 Farms	*		Statement A is incorrect
	(2)	P3 Farms		(3)	Both Statement A and Statement B are incorrect
	(3)	P2 Farms		(4)	Both Statement A and Statement B
	(4)	P1 Farms			are correct
545		(3-	S)		

FC

6. Consider the following statements with respect to mulberry:

Statement A: The flowers of mulberry are unisexual.

Statement B: The cross-pollination takes place in mulberry.

Statement C: The chief pollinating agent in mulberry is wind.

Choose the correct answer using the codes below:

- (1) A and B only
- (2) B and C only
- (3) A, B and C
- (4) A and C only
- 7. Which among the following is/are wrong statements?

Statement A: The female silkworm pupa are heavier than males.

Statement B: Females have tapering abdomen with x mark.

Statement C: Male silk moths are more active than female.

Statement D: Female moths emerge earlier than males in Mysore seed cocoons.

- (1) A only
- (2) A and B only
- (3) B and D only
- (4) A and C only

- 8. With reference to the food plants of non-mulberry silkworms, consider the following statements:
 - A. The Tasar silkworm feeds on the leaves of *Terminalia arjuna* and also the *Terminalia tomentosa*.
 - B. The Muga silkworm feeds on only the leaves of Machilus bombycina.
 - C. The Eri silkworm feeds on the leaves of Ricinus communis but does not feed on the leaves of Manihot utilissima.

Which of the above statements is/are correct?

- (1) A only
- (2) B only
- (3) Conly
- (4) B and C only
- 9. High proportion of carotenoids is found in the cocoon of:
 - (1) Indian race
 - (2) European race
 - (3) Korean race
 - (4) Indo-China race
- 10. Pirn winding is necessary to prepare:
 - (1) Warp
 - (2) Skein
 - (3) Weft
 - (4) All of the above

- 11. With reference to the spread of sericulture to India, consider the following statements:
 - A. The silk industry is said to have spread to Tibet when a Chinese princess, carrying silkworm eggs in her head-dress, married the king of Khotan in Tibet.
 - B. China conquered Khotan in Tibet and introduced sericulture.
 - C. From Tibet, the sericulture industry spread to India.

- (1) A and B only
- (2) B and C only
- (3) A and C only.
- (4) A, B and C
- 12. With reference to the sericulture development by the British East India Company, consider the following statements:
 - A. When the British came to India, they found a flourishing silk trade.
 - B. The East India Company exploited the industry and developed silk centres in India. In 1836 because of the trade being turned over to the private enterprises, the silk industry declined in West Bengal.
 - C. This resulted in decline of sericulture also in Mysore, Jammu and Kashmir.

Which of the above statements are correct?

- (1) A and B only
- (2) B and C only
- (3) A and C only
- (4) A, B and C

- of the Britishers export large quantities of raw silk to England?
 - (1) Jammu and Kashmir
 - (2) Karnataka
 - (3) Tamil Nadu
 - (4) West Bengal
 - 14. With reference to the sericulture practice in different climates, consider the following statements:
 - A. Sericulture is practised in both tropical and temperate climate conditions.
 - B. However, in temperate climate the cultivation of mulberry is acclimatized, which ensures continuous supply of mulberry leaf for silkworm rearing.
 - C. Whereas, in tropical climate due to adequate climatic conditions, suitable soil, good sunshine and rainfall, the quantity of the silk production is more, which sustains the sericulture industry.

Which of the above statements are correct?

- (1) A and B only
- (2) B and C only
- (3) A and C only
- (4) A, B and C

15. Consider the following pairs:

Textile Fibre

Origin

- A. Viscose
- Regenerated
- B. Ramie
- Synthetic
- C. Cotton
- Plant
- D. Polypropylene -
- Regenerated

How many pairs given above are correctly matched?

- (1) Only one pair
- (2) Only two pairs
- (3) Only three pairs
- (4) All four pairs
- 16. Sericulture from China and India was carried to Europe along the famous Silk Road. The correct order in which this road passed through the following cities is:
 - (1) Baghdad, Tashkent, Damascus and Istanbul
 - (2) Tashkent, Baghdad, Damascus and Istanbul
 - (3) Damascus, Baghdad, Tashkent and Istanbul
 - (4) Istanbul, Damascus, Baghdad and Tashkent

- 17. Mulberry plant is considered as:
 - Perennial, Dicot, Dioecious and Bisexual
 - (2) Perennial, Dicot, Monoecious and Bisexual
 - (3) Perennial, Monocot, Dioecious and Bisexual
 - (4) Perennial, Dicot, Dioecious and Unisexual
- 18. Consider the constituents of the mulberry fruit:

	Constituent		Percent (%)
A.	Moisture	I.	1.5
В.	Protein	11.	8.3
0-	Carbohydrate	Ш.	87.5

- D. Fat
- IV. 0.5

V. 0.4

Choose the correct answer from the options given below:

	A	В	\mathbf{C}	D
(1)	I	IV	V	Ш
(2)	V	II	III	I
(3)	II	III	V	ſV
(4)	III	I	II	V

- 19. The commercial propagation of mulberry is through:
 - (1) Graftings
 - (2) Seedlings
 - (3) Cuttings
 - (4) Layerings



Website www.shikarat.com

For more contact us



Instagram



Email

Admin@shikarat.com









Twitter



Sharing is Caring





fit is helpful Please **Share with your** friends 😃

20.	Consider	the	following	mutagenic
	chemicals:			72

- A. Ethyl Methane Sulphonate
- B. Methyl Methane Sulphonate
- C. 1,4-Dimethyl Sulphonoxy butane
- D. 5-Bromouracil

Which of the above are used in mutation breeding of mulberry?

- (1) A and B only
- (2) B and C only
- (3) A and C only
- (4) A, B, C and D

 ♣

21. Consider the mulberry varieties:

Variety

Source

- A. V1
- I. RSRS, Coonoor
- B. Viswa (DD) II. KSSRDI, Thalaghattapura
- C. MR2 III. Kanva Silk Farm, Channapatna
- D. M5 IV. CSRTI, Mysuru

Choose the correct answer from the options given below:

	A	В	\mathbf{C}	D	
(1)	IV	II	I	III	
(2)	I	III	II	IV	
(3)	III	IV	II	I	
(4)	III	I	IV	II	

- 22. The term 'Emasculation' refers to:
 - (1) Collection of stamens from male catkin
 - (2) Collection of pollen from male catkin
 - (3) Collection of female flowers from female catkin
 - (4) Collection of male flowers from male catkin
- 23. The suitable soil pH for mulberry cultivation is:
 - (1) 5.4 5.8
 - (2) 6.4 6.8
 - $(3) \quad 6.8 7.8$
 - $(4) \quad 7.8 8.8$
- 24. Consider the following pairs:

Mulberry Nativity Species/Varieties

- A. Morus indica West Java
- B. Gelso romano Europe -
- C. Morus alba L. India
- D. Morus nigra South Vietnam

How many pairs given above are correctly matched?

- (1) Only one pair 🗡
- (2) Only two pairs
- (3) Only three pairs
- (4) All four pairs ×

(11 - S)

25.	'Seriboost'	in	mulberry	is	used	as	
			**********		CALLE	440	

- Chemical fertiliser (1)
- Organic manure (2)
- (3) Green manure
- Micronutrient (4)
- 26. With reference to the application of fertiliser/manure in irrigated mulberry consider the following cultivation. statements:
 - Farmyard Manure (FYM) should be A. applied @ 20_tonnes/ha/year as recommended for irrigated mulberry.
 - Azotobacter Apply 20 kg of B. biofertiliser/ha/year in 5 split doses @ 4 kg each time after every leaf harvest/priming and inter-cultivation operations.
 - It has also been recommended that C. application of 300 kg urea along with 20 kg Azotobacter helps in good growth and yield of mulberry.

- A and B only (1)
- (2)B and C only
- (3)A and C only
- A, B and C (4)



The percentage of Nitrogen present in Farmyard Manure (FYM) is:

- 0.01 0.02%
- 0.02 0.03%(2)
- (3)
- (4)
- 0.03 0.04%0.3 - 0.5%
- Considering the diseases of mulberry, 28. match List I with List II:

List 1

List II

- A. Mulberry leaf blight
- I. Helicobasidium mompa
- II. Pseudomonas mori Violet root
- Mulberry III. Cerotelium fici C. leaf spot
- Mulberry IV. Cercospora moricola leaf rust
 - V. Rosellinia necatrix

Choose the correct answer from the options given below:

I

II

	A	В	\mathbf{C}	D
(1)	II	Ι	IV	III
(2)	I	II	III	IV
(3)	IV	V	II	III

III

(4)

 Γ V

- With reference to the powdery mildew 29. disease in mulberry leaf, consider the following statements:
 - Asexual reproduction takes place by A. the means of fruiting bodies called cleistothecia which give rise to conidia.
 - During favourable conditions, the B. Phyllactinia corylea pathogen through germination, develops mycelia, conidiophores and conditions which lead to the spread of disease.
 - C. The Phyllactinia corylea multiplies the host only by asexual reproduction reducing the quality.

- (1)A and B only
- B and C only (2)
- **(3)** A and C only
- B only (4)

Consider the following pairs: 30.

> Spores Disease

- Mulberry leaf rust Basidiospores
- White root rot Ascospores В.
- Uredospores C. Violet root rot
- Mulberry leaf spot Conidiophores How many pairs given above are correctly matched?
- Only one pair (1)
- Only two pairs (2)
- (3)Only three pairs
- All four pairs (4)
- Match List I with List II: 31.

List II List I

- Mulberry I. Sthenias Grasshopper grisator
- Mealy bugs II. Spilosoma obliqua
- Bihar hairy III. Maconellicoccus caterpillar hirsutus
- Mulberry leaf IV. Neorthacris roller acuticeps nilgiriensis
 - V. Morgaronia pulverulentalis

Choose the correct answer from the options given below:

	A	В	C	D
(1)	V	IV	III	II
(2)	IV	Ш	II	V
(3)	II	I	IV	III
(4)	III	IV	v	II

32.	8th	abdom	inal seg	ment	dal horn' on of the silkw f the order :		35.		presence of 'the Ishiwata's glands' in vorm larvae are considered to be the
	(1)	Hemi	ptera					genit	al markings on the ventral side of :
	(2)	Ortho	optera					(1)	The 8th segment of male larva
	(3)	Lepid	loptera					(0)	Mr. oth segment of female large
	(4)	Thysa	anoptera	l.				(2)	The 8th segment of female larva
33.			ntal cro lar Gold'		the silkw	vorm		(3)	The 8 th and 9 th segments of female larva
	(1)	d CS	R-2 × PN	1º				(4)	The 8th and 9th segments of male
	(2)	d CS	R-4 × HI	ΜŶ					larva
	(3)	d CS	R-2 × Ni	rtary (?				all a
	(4)	d CS	$R-4 \times C$.	Nichi	۹ >		36.	The	function of the 'maxillae' in silkworm
34.	silk	worm la	_		fication of arts, match L	the List I		larva	
	with	h List II	:		7		2	(1)	Photoreceptor
		List I Body S	legment		List II Mouth Parts	s	7	(2)	Mechanoreceptor
	A.	Second		I.	Labium	7,		(3)	Chemoreceptor
	B.	Fourth		II.	Mandibles			(4)	Thermoreceptor
	C.	Fifth		III.	Antennae				,
	D.	Sixth		Ly.	Maxilla		37.	The	enzyme secreted by the silk moths of
			correct		swer from	the		Bom	byx mori is:
		A	В	C	D			(1)	Fibrinase
((1)	III	II	IV	I			(2)	Sericinase
	2)	I	III	II	IV	i			
(3)	IV	II	I	III			(3)	Amylase
(-	4)	II	IV	III	I			(4)	Cocoonase
545						(17 -	- S),		

- 38. The sex attractant pheromone is secreted in the silkworm *Bombyx mori* to ensure the mating from which part of the following?
 - (1) Labin of the female silk moth
 - (2) Harper of the male silk moth
 - (3) Sacculi lateralis of the female silk moth
 - (4) Ovipositor of the female silk moth
- 39. With reference to the anatomy of silkworm Bombyx mori, consider the following:

Anatomical Organs system

- A. Digestive Cardiac valve system
- B. Circulatory Malpighian y system tubules
- C. Respiratory Tracheoles / system
- D. Excretory Ostia

How many pairs given above are correctly matched?

- (1) Only one pair
- (2) Only two pairs
- (3) Only three pairs
- (4) All four pairs

- 40. The juveni<u>le horm</u>one in silkworm

 Bombyx mori is secreted from:
 - (1) The neurosecretory cells of brain
 - (2) Corpora cardiaca
 - (3) Corpora allata
 - (4) Prothoracic glands
- 41. With reference to the reproductive system of female silk moth, match the following:

List I List II

- A. Ovarioles I. Adhesive substance
- B. Accessory II. Storage of Reproductive sperms glands
- C. Spermatheca III. Synthesis of eggs
- D. Bursa IV. Receiving of sperms

 V. Receiving of eggs

Choose the correct answer from the options given below:

	A	В	C	D	
(1)	I	IV	V	III	
(2)	V	II	III	I	
(3)	III	I	II	IV	
(4)	II	III	I	V	

- **42.** Which part of the silk glands secretes Sericin III?
 - (1) Posterior region of the silk glands
 - (2) Middle region of the silk glands
 - (3) Posterior part of the middle region of the silk glands
 - (4) Anterior part of the middle region of the silk glands
- 43. Consider the following statements:
 - A. The sperm is received into the female silk moth and makes internal fertilization.
 - B. The sperm is received into the female silk moth and is stored in the spermatheca.
 - C. The stored sperm enters into the egg and makes external fertilization.

- (1) A and B only
- (2) A and C only
- (3) B and C only
- (4) A, B and C

- 44. How many times does the silkworm increase in its weight from the day of hatching to maturity, i.e., last day of fifth instar?
 - (1) 7,000 times
 - (2) 10,000 times
 - (3) 6,000 times
 - (4) 5,000 times
- 45. The silkworm larva is blind due to:
 - (1) Lack of ocelli
 - (2) Lack of eyes
 - (3) Lack of compound eyes ⊁
 - (4) Lack of sense organs
- 46. The cells which secrete the digestive enzymes in silkworm larva are:
 - (1) The cells of Foregut
 - (2) The Goblet cells
 - (3) The Cylindrical cells
 - (4) The cells of Hindgut

(21 - S)

47.	are	the br	nt of sill ain, sub	worm ooesop	the embryonic , Bombyx mori, hageal ganglion formed?		BmDNV-1 infection is determined by a recessive gene called: (1) Nid-1
	(1) (2)			- 2	ng of the egg		(2) nsd-1 (3) nsd-Z (4) Reg-1
	(3)		to the state of th		g of the egg	50.	The number of the endocrine glands in silkworm are:
48.	(4)		following		ing of the egg		(1) 5 (2) 3 (3) 4
	A.	Tasar I	Ecoraces	Ī.	Food Plants Zizyphus	51.	(4) 6 Consider the following pairs in Bombyx
	B.	Jiriban	n	II.	Hardwickia Terminalia	F	mori : Sense Organs Type of Sensation
•	C. D.	Raily Belgau	m		Shorea		A. Antennae – Mechanoreceptor > B. Integument – Photoreceptor
			e correc	11.	swer from the		C. Maxillae – Chemoreceptor / D. Compound – Thermoreceptor /
	(1)	A _.	В	C	D By		eyes How many pairs given above are correctly
×	(1)	II	I	II IV	IV III		matched? (1) Only one pair
	(3)	III	I	IV	п		(2) Only two pairs(3) Only three pairs
	(4)	IV	III	II	I		(4) All four pairs
545					(23	3-S))

Match the following: 52.

Name of the Silkworm

Chromosome Number

- Bombyx mori
- I. n = 28
- B. Bombyx mandarina
- II. n = 31
- C. Antheraea mylitta
- III. n = 27
- D. Philosamia ricini
- IV. n = 15

V. n = 14

Choose the correct answer from the options given below:

> В A

- C \mathbf{D} .
- I II (1)
- П
- IIIIV
- (2)III
- IV

II

I

- I (3)
- III
- IV (4)

- the primary and On which day are **53.** secondary spermatogonia differentiated in the silkworm Bombyx mori?
 - Three days after hatching (1) 70
 - Four days after hatching (2)
 - Six days after hatching (3)
 - Eight days after hatching (4)

- 54. The genes responsible for moulting in Bombyx mori are:
 - V-alleles (1)
 - M-alleles (2)
 - Lm-alleles (3)
 - Autosomes
- In silkworm Bombyx mori, the female 55. sex-determining factor is present in:
 - 'Z' chromosome (1)
 - 'W' chromosome (2)
 - 10th Autosome
 - 12th Autosome
- Consider the following pairs:

Ectodermal Silkworm Stage Derivative

- Larva
- Integument
- В. Prepupa
- Scales
- Pupa C.
- Chitin
- D. Adult moth
- Chitin

How many pairs given above are correctly matched?

- Only one pair (1)
- Only two pairs (2)
- (3)Only three pairs
- (4) All four pairs

- 57. The thoracic ganglion in the silkworm larva gives rise to:
 - (1) One pair of nerves
 - (2) Two pairs of nerves
 - (3) Three pairs of nerves
 - (4) Four pairs of nerves
- 58. The Japanese scientists produced parthenogenetic individuals by treating the unfertilized eggs of silkworm through:
 - (1) HCl with specific gravity 1.04 1.06 at $70 74^{\circ}$ F for 4 6 minutes
 - (2) HCl with specific gravity 1.06 1.08 at $70 74^{\circ}$ F for 8 10 minutes 9
 - (3). HCl with specific gravity 1.08 1.10 at $70 74^{\circ}$ F for 4 6 minutes
 - (4) HCl with specific gravity $1.075 1.10 \text{ at } 70 74^{\circ} \text{ F for } 4 5$ minutes

545

59. With reference to the sex-limited races in silkworm, consider the following pairs:

State Character

- A. Male larva Presence of dark
 markings in
 thorax
- B. Female larva Plain and no markings in thorax
- C. Pupa No markings 🗴
- D. Cocoon The females are coloured and males are plain

How many pairs given above are correctly matched?

- (1) Only one pair
- (2) Only two pairs
- (3) Only three pairs
- (4) All four pairs

(27 - S)

- **60.** With reference to silkworm breeding, consider the following statements:
 - A. The polyvoltine races are unique characteristic features with less larval duration and resistant to diseases considered as female parent to cross with male parent of bivoltine.
 - B. However, many of the polyvoltine races may not have good combining ability with that of bivoltine races.
 - C. For the production of F_1 Hybrids the female parent of bivoltine is crossed with the male parent of polyvoltine. The F_1 hybrids of such cross are considered for commercial silk production. (Example: $\P NB4D2 \times PM \circlearrowleft$) \swarrow

- (1) A and B only
- (2) B and C only
- (3) A and C only
- (4) A, B and C

- 61. Which of the following breed is evolved by CSR & TI Mysore?
 - (1) Shunrei
 - (2) Kinshu
 - (3) NB₇
 - (4) KA
- **62.** With reference to the production of hybrids in silkworms, consider the following:
 - A. $FC_1 \times FC_2$ Double hybrids –
 - B. $PM \times NB_4O_2$ Multivoltine hybrid
 - C. CSR-1 × CSR-2 Cross breed
 - D. $PM \times FC_1$ Poly hybrid

How many pairs given above are correctly matched?

- (1) Only one pair
- (2) Only two pairs
- (3) Only three pairs
- (4) All four pairs

- 63. The pure races of silkworms have higher significance in the production of cross breeds. Consider the following statements:
 - A. The pure stocks of multivoltine and bivoltine races are reared giving maximum care with recommended rearing technology.
 - B. Maintenance of the racial characters of such races are so important which may reflect in their F₁ hybrids to get higher qualitative and quantitative characters.
 - C. The rearing factors may not be the major criteria to get Heterosis in the cross breeds.

Which among the above statements is/are incorrect?

- (1) A only
- (2) B only
- (3) C only
- (4) A, B and C
- 64. The 'Black boxing' process for the silkworm eggs is done to ensure:
 - (1) early hatching.
 - (2) delay in hatching.
 - (3) difference in hatching.
 - (4) uniform hatching.

- 65. In the silkworm egg production centres (Grainages), consider the following factors related to Oviposition:
 - A. The success of Oviposition largely depends on the mating duration of the silk moths. Normally, mating is allowed for 3 hours to ensure two to three ejaculations essential for fertilization of eggs.
 - B. The mating of male silk moths for more than 2 times with virgin females does not matter for maximum fecundity and fertilized eggs.
 - C. During oviposition, the de-coupled female silk moths are maintained in a semi-dark condition preferably at a temperature and humidity of 25 ± 1°C and 75 ± 5% respectively.

Which of the above factors is/are correct?

- (1) A only
- (2) A and B only
- (3) A and C only
- (4) A, B and C

- 66. The National Silkworm Seed Project (NSSP) has developed the silkworm seed multiplication programme for the supply of quality layings. Choose the correct sequence of the following:
 - (1) $P1 \rightarrow P2 \rightarrow P3 \rightarrow P4$
 - $(2) \quad P4 \rightarrow P2 \rightarrow P3 \rightarrow P1$
 - $(3) \quad P4 \rightarrow P3 \rightarrow P2 \rightarrow P1$
 - $(4) \quad P3 \rightarrow P2 \rightarrow P1$
- 67. The orientation of the rearing house in tropical zones should be:
 - (1) North South
 - (2) East West
 - (3) North East
 - (4) South East
- 68. With reference to the disinfectants in silkworm rearing, consider the following pairs:
 - A. Sanitech Bed disinfectant
 - B. Bleaching Room solution disinfectant
 - C. Vijetha Bed disinfectant –
 - D. Sanjeevani Room disinfectant

How many pairs given above are correctly matched?

- (1) Only one pair
- (2) Only two pairs
- (3) Only three pairs
- (4) All four pairs

- 69. With reference to the 'chawki rearing' consider the following statements:
 - A. The chawki worms grow extremely fast under high temperature (up to 30°C) and high humidity (up to 90%), good ventilation and fed with tender nutritious leaves.
 - B. Even though the consumption of mulberry leaf is only 6% of the total feed but shows 400 times increase in body weight.
 - C. For success of chawki rearing, quality of leaves and maintenance of ideal temperature and humidity plays a very significant role.

Which of the above statements are correct?

- (1) A and B only
- (2) B and C only
- (3) A and C only
- (4) A, B and C

With reference to the temperature and 72. Which among the following is considered 70. relative humidity in silkworm rearing, consider the following pairs:

Age	Temperature	Relative
		Humidity

- II Instar $27 - 28^{\circ}C$ 85 - 90%
- III Instar В. $22 - 23^{\circ}C$ 65 - 70% >
- C. **IV** Instar $25 - 26^{\circ}C$ 70 - 75%
- D. V Instar $27 - 28^{\circ}C$ 85 - 90%

How many pairs given above are correctly matched?

- (1)Only one pair
- (2)Only two pairs
- (3)Only three pairs
- (4) All four pairs
- The 71. characteristic features moulting larva are:
 - (1) Stops feeding, inactive, becomes stout and shiny and head becomes dark.
 - Stops feeding, active, loose skin and (2) head becomes swollen. >
 - Stops feeding, less active, elasticity (3) of the body and head becomes transparent.
 - Stops feeding, immobile, loose skin (4) and head becomes grey in colour.

- as ideal cocoon spinning equipment?
 - Paddy straw mountage > (1)
 - Bottle brush mountage (2)
 - Bamboo mountage Chandrika (3)
 - Rotary mountage (4)
- The ideal time of cocoon harvesting for 73. temperate silkworm races is:
 - 3rd or 4th day after spinning (1)
 - 4th or 5th day after spinning (2)
 - 5th or 6th day after spinning (3)
 - 7th or 8th day after spinning
- With reference to the development of Nosema bombycis consider the following:
 - A. Mature spore - Meront - Planont -Amoebula
 - Mature spore Planont Amoebula В. - Meront
 - C. Mature spore - Amoebula - Meront - Planont
 - D. Mature spore - Amoebula - Planont Meront.

Which of the above sequence is correct?

- (1)A
- **(2)** B
- (3) C
- (4) D

75.	With reference to the site of infection of the pathogen in silkworm diseases, match the following:					
		Pathogen		Site of infection		
	A.	NPV	I.	The cytoplasm of cylindrical cells of		

- midgut
- B. CPV II. Nuclei of the columnar cells of midgut
- C. IFV III. Nuclei of the tracheal epithelial cells
- D. DNV IV. Cytoplasm of the goblet cells of midgut

Choose the correct answer from the options given below:

	A	В	\mathbf{C}	D	
(1)	I	II	III	IV	
(2)	IV	III	II	I	
(3)	III	I	IV	II	
(4)	II	IV	I	III	

Consider the following:

	Name		Family
A.	Nasolynx thymus	_	Eulophidae
B.	Trichopria species	4	Chalcidae
C.	Exoristobia philippinensis	5	Encyrtidae
D.	Spilomicrus karnatakensis	-	Diapriidae 🥦

How many pairs given above are correctly matched?

- (1) Only one pair
- (2)Only two pairs
- (3)Only three pairs
- (4) All four pairs

- 77. Among the silkworm diseases, there is a disease called "no by o" in Japanese which is actually called:
 - Flacherie (1)
 - Grasserie (2)
 - (3)Muscardine
 - Pebrine (4)
- With reference to the silkworm diseases, 78. match the following:

1	Disease		Causative Agent
87	Nuclear polyhedrosis	I.	Serratia marcescens
В.	Red septicemia	II.	Beauveria bassiana
C.	White muscardine	III.	Nosema bombycis
D.	Pebrine	IV.	Borrelina bombycis

Choose the correct answer from the options given below:

	A	В	C	D
(1)	I	II	III	ΙV
(2)	II	III	IV	I
(3)	IV	I	II	III
(4)	III	IV	I	II

- 79. The silkworm infected with white muscardine, unlike other diseases does not rot or decay but remains hard because:
 - the fungus secretes crystals of ammonium and magnesium.
 - (2) the fungus secretes crystals of ammonium and calcium.
 - (3) the fungus secretes magnesium and calcium.
 - (4) the fungus secretes only the crystals of ammonium.

- 80. Which among the following is not considered for the biological control of uzi fly?
 - (1) Nesolynx thymus
 - (2) Trichopria species
 - (3) Exoristobia philippinensis
 - (4) Trichogramma chilonis

- 81. In which stage of the silkworm Bombyx mori is the Dermestid beetle infested?
 - (1) Egg
 - (2) Larva
 - (3) Cocoon
 - (4) Moth
- 82. With reference to the sericulture extension, consider the following statements:
 - A. The CSB and State sericulture organisations are giving maximum effort for improvisation in mulberry and cocoon production.
 - B. Creating awareness and transfer of innovative technology into the field is the prime duty of the sericulture extension personnel.
 - C. Apart from farm and home visit by the extension workers, other means of transferring the innovative knowledge may not be needed. >

- (1) A and B only
- (2) B and C only
- (3) A and C only
- (4) A, B and C

(39 - S)

- g3. With reference to the transaction of cocoons in the cocoon markets, consider the following statements:
 - A. The cocoon transaction is carried out in the Government prescribed cocoon markets only.
 - B. Only authorised cocoon rearers and reelers are allowed to participate in cocoon transactions. The process is by open auction and e-auction in the presence of marketing officials.
 - C. In certain circumstances, the cocoon transaction by the reelers may not be limited beyond the amount deposited by him.

- (1) A and B only
- (2) B and C only
- (3) A and C only
- (4) A, B and C

84. Expand KSMB:

- (1) Karnataka Silk Mobilizing Board
- (2) Karnataka Silk Motivating Board
- (3) Karnataka Silk Marketing Board
- (4) Karnataka Silk Modernising Board

85. Given below are two statements with respect to silk testing:

Statement A: Elongation is the amount of stretch when the yarn is pulled to its breaking point.

Statement B: The apparatus used for tenacity test is serigraph.

In light of above statements, choose the most appropriate answer.

- (1) Statement A is correct but Statement B is incorrect
- (2) Statement B is correct but Statement A is incorrect
- (3) Both Statement A and Statement B are incorrect
- (4) Both Statement A and Statement B are correct
- 86. Which of the following is **not** a commercial method of stifling?
 - (1) Sun drying
 - (2) Hot air drying
 - (3) Steam stifling
 - (4) Cold air killing

- With reference to the steam stifling 90. With reference to the silk reeling 87. consider the following statements :
 - A. Steam stifling not only kills the pupae of the cocoon, it also ensures softening of the sericin.
 - Thus, it facilitates good cocoon B. cooking, helps in easy unwinding of silk filaments at the time of reeling.
 - The steam stifled cocoons can be C. preserved for long durations up to 6 months.

- (1) A only
- (2)B only
- (3)A and B only
- (4) A and C only
- The pH of the reeling water is: 88. MNS
 - (1) 6.0 - 6.2
 - (2) 6.2 - 6.4
 - (3) 6.4 - 6.6
 - (4) 6.8 - 8.4

545

- The 'Rendita is a deciding factor of: 89.
 - Less cocoon quantity with higher (1)silk production.
 - Higher cocoon quantity with higher (2)silk production.
 - Equal cocoon quantity with equal (3)silk production.
 - Higher cocoon quantity with lower (4) silk production.

- machine, which among the following have higher efficiency in silk reeling process?
 - Improved Charaka (1)
 - Cottage Basin (2)
 - (3)Charaka
 - Multi-end-reeling machine (4)
- The circumference of the small reels of 91. multi-end-reeling machine is:
 - 60 61 cm(1)
 - 65 70 cm(2)
 - 70 80 cm
 - (4) 80 85 cm
- With reference to the reeling devices in reeling machines, consider the following pairs:

Reeling Function Devices

- Jettebout Feeding/casting of silk filaments
- В. Button Feeding of silk filaments
- C. Traverse Passing silk thread through hole
- D. Reel Winding of silk filaments

How many pairs given above are correctly matched?

- (1)Only one pair
- (2)Only two pairs
- (3)Only three pairs
- (4) All four pairs

93.	Whicond	ch of the following silk tests are ucted in Seriplane machine?	98.		relevar mulber
	(1)	Evenness variation test			wing:
	(2)	Tenacity and elongation test			Equipn
	(3)	Winding test			
	(4)	Cohesion test		A.	Chalon
94.	The	correct sequence of the silk throwing			Khora
	(1)	Reeling \rightarrow Doubling \rightarrow Twisting \rightarrow Re-winding			Khorik. Jali
	(2)	Winding \rightarrow Doubling \rightarrow Twisting \rightarrow Re-winding		D . (Jan
	(3)	Winding \rightarrow Re-winding \rightarrow Doubling \rightarrow Twisting		Choc	ose the
	(4) •	$\begin{array}{c} \text{Doubling} \rightarrow \text{Winding} \rightarrow \text{Twisting} \rightarrow \\ \text{Re-winding} \end{array}$		(1)	AC)
95.	The	widthwise/horizontal threads in the		(2)	v
	fabri	c are:	0	(3)	IV
	(1)	Warps	LY	(4)	III
	(2)	Wefts	99.	W/L:	.b
	(3)	Pirn	99.		ch amo
	(4)	Shuttle			Techn
96.	The	liquor ratio in silk dyeing process is :		(1)	Nosen
00.	(1)	1:20		(2)	NPV
	(2)	1:30			
	(3)	1:40		(3)	Beauv
	(4)	1:50		(4)	Asper
97,		parentage of FCI Bivoltine silkworm id is:	100.	The from	indige:
	(1)	$CSR 6 \times CSR 26$		(1)	West
	(2)	$CSR\ 2 \times CSR\ 27$		(2)	Assan
		BL $67 \times CSR$ 101		(3)	Karna
	(3)			(4)	Jamm
	(1)	$CSR 2 \times CSR 4$	I	,	Guinn

98.	In	relevance				uipment	
	non-mulberry		silkworms,		s,	, match	the
	foll	owing:					

	Equipment		Function
A.	Chaloni	I.	Spinning equipment
В.	Khora	II.	Coupling of male and female moths
\mathbf{C} .	Khorika	III.	Collection of ripened worms
D.	Jali	IV.	Transferring of worms
Ch	aga the co	V.	Moth cage

en below:

	Α	В	C	D
(1)	$\widetilde{}$	II	III	ΙV
(2)	V	I	IV	III
(3)	IV	III	II	I
(4)	III	V	IV	II

- ng the following, in silkworm, ed as plasmid in Recombinant ology?
 - na bombycis
 - eria bassiana
 - gillus flavus
- nous silkworm race Moria is
 - Bengal

 - ataka
 - nu and Kashmir